



TITLE:

DROPLETS FROM THE PLANKTON NET XXIV. THE PREDOMINANT TYPE OF VELELLA IN JAPANESE WATERS

AUTHOR(S):

Bieri, Robert

CITATION:

Bieri, Robert. DROPLETS FROM THE PLANKTON NET XXIV. THE PREDOMINANT TYPE OF VELELLA IN JAPANESE WATERS. PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY 1970, 17(5): 303-304

ISSUE DATE:

1970-02-21

URL:

<http://hdl.handle.net/2433/175608>

RIGHT:

DROPLETS FROM THE PLANKTON NET XXIV. THE PREDOMINANT TYPE OF *VELELLA* IN JAPANESE WATERS¹⁾

ROBERT BIERI

Antioch College, Yellow Springs, Ohio 45387

With 1 Table

The appearance of *Katsuonokanmuri* or bonito's crown along the coast of Japan heralds the approach of warm water and good fishing for bonito. How fine that *Velella*, bringing such good news, also adds only beauty to the sea scape—no poisonous stings!

From November 1965 to June 1966, I recorded the occurrence of *Velella* along the beaches of Shirahama. A summary of my data is given in Table 1. *Velella*

Table 1. Records of *Velella* strandings, Shirahama

<i>Date</i>	<i>Type</i>	<i>Number</i>	<i>Size Range mm</i>
Nov. 5, 1965	Left	5	12 -25
Nov. 8, 1965	Left	15	9 -30
Nov. 9, 1965	Left	31	9 -38
	Right	6	3 -18
Mar. 4, 1966	Left	230	5.5-50
Mar. 5, 1966	Left	51	15 -58
Mar. 23-24, 1966	Left	70	14 -36
	Right	1	22
April 27, 1966	Left	283	7 -62
	Right	80	17 -61
June 8, 1966	Left	5	16 -29

Total: Left 690, Right 87

was stranded only nine days out of 230. In all, with the generous help of Mr. H. TANASE and some sailors of the glass bottom boats, I collected 777 specimens of *Velella*. Six-hundred ninety or 90% of them were left handed; that is, if the animal is held with the long axis north and south the crest runs from northwest to southeast. The remainder were right handed. In *Velella*, left handers sail to the left of the wind.

1) Contributions from the Seto Marine Biological Laboratory, No. 515.

Thus the theory of SAVILOV (1958–1961) is upheld. Left handed *Velella* sailing to the left of the wind should be most abundant along the outer edges of the clock-wise rotating wind and current system. The right-handers should be most abundant inside such a gyre. But winds and currents are variable with many eddies; so it is to be expected that nature does not perfectly fulfill our theories. In this case, she contradicts our predictions 10% of the time.

REFERENCES

- SAVILOV, A.I. 1958. Pleuston of the Western Part of the Pacific Ocean. Doklady Akad. Nauk SSSR, Vol. 122, No. 6, pp. 1010–1017. (in Russian)
- 1961. The Distribution of the Ecological Forms of the By-the-Wind-Sailor, *Velella lata* CH. and Eys., and the Portuguese Man-of-War, *Physalia utriculus* (LA MARTINIERE) ESCH., in the North Pacific. Trudy Inst. Okeanol. Akad. Nauk SSSR, Vol. 45, pp. 223–239. (in Russian)